

Nursing Home Compare - Measure Dates

Measure Code
401
402
403
404
405
406
407
408
409
410
411
415
419
424
425
426
430
434

Nursing Home Compare - Measure Dates

Measure Description

Percent of long-stay residents whose need for help with daily activities has increased. Lower percentages are better.

Percent of long-stay residents who self-report moderate to severe pain. Lower percentages are better.

Percent of long-stay high-risk residents with pressure ulcers. Lower percentages are better.

Percent of long-stay residents who lose too much weight. Lower percentages are better.

Percent of long-stay low-risk residents who lose control of their bowels or bladder. Lower percentages are better.

Percent of long-stay residents who have/had a catheter inserted and left in their bladder. Lower percentages are better.

Percent of long-stay residents with a urinary tract infection. Lower percentages are better.

Percent of long-stay residents who have depressive symptoms. Lower percentages are better.

Percent of long-stay residents who were physically restrained. Lower percentages are better.

Percent of long-stay residents experiencing one or more falls with major injury. Lower percentages are better.

Percent of long-stay residents assessed and given, appropriately, the seasonal influenza vaccine. Higher percentages are better.

Percent of long-stay residents assessed and given, appropriately, the pneumococcal vaccine. Higher percentages are better.

Percent of long-stay residents who received an antipsychotic medication. Lower percentages are better.

Percent of short-stay residents who self-report moderate to severe pain. Lower percentages are better.

Percent of short-stay residents with Pressure ulcers that are new or worsened. Lower percentages are better.

Percent of short-stay residents assessed and given, appropriately, the seasonal influenza vaccine. Higher percentages are better.

Percent of short-stay residents assessed and given, appropriately, the pneumococcal vaccine. Higher percentages are better.

Percent of short-stay residents who newly received an antipsychotic medication. Lower percentages are better.

Nursing Home Compare - Measure Dates

Start Date	End Date
10/01/2011	06/30/2012
10/01/2011	06/30/2012
10/01/2011	06/30/2012
10/01/2011	06/30/2012
10/01/2011	06/30/2012
10/01/2011	06/30/2012
10/01/2011	06/30/2012
10/01/2011	06/30/2012
10/01/2011	06/30/2012
10/01/2011	06/30/2012
10/01/2011	06/30/2012
10/01/2011	06/30/2012
10/01/2011	06/30/2012
07/01/2011	06/30/2012
07/01/2011	06/30/2012
07/01/2011	06/30/2012
07/01/2011	06/30/2012
07/01/2011	06/30/2012